

APPROVED CPE

NOW ONLINE: www.acp.edu.au or www.jnlcompmed.com.au

CPE article 21: calendula

Pharmacists This unit is worth half-an-hour CPE credit. ACPPM members and CPE self-assessment program subscribers can read the Calendula monograph, then circle the appropriate answers and complete the form below the questions, using a black pen. Photocopy the page and fax it to the Australian College of Pharmacy Practice and Management on 02 6273 8988, or post to: ACPPM, PO Box 7007, CANBERRA BC ACT 2610. Or, complete the unit online: www.acp.edu.au. Be prompt — ACPPM must receive your answers by 2 January 2006. Answers will be printed in the next JCM [Vol 5(1)] in January/February 2006.

Q Circle the most appropriate answer to the following questions. More than one answer may be correct.

- 1 Calendula has been used traditionally for
 - a sunstroke
 - b burns
 - c snake bite
 - d poor wound healing
- 2 The main pharmacological actions of calendula are
 - a anti-inflammatory, lipid lowering
 - b anti-inflammatory, hypoglycaemic
 - c promotion of wound healing, anti-inflammatory
 - d antidepressive, antimicrobial
- 3 According to a randomised controlled study conducted with breast-cancer patients
 - a a calendula tincture taken orally prevents radiationinduced dermatitis
 - b a calendula-flower preparation used topically prevents radiation-induced dermatitis
 - c a calendula-flower preparation used topically is ineffective for preventing radiation-induced dermatitis
 - d a calendula tincture applied topically is ineffective for preventing radiation-induced dermatitis

- 4 According to the same study, topical calendula treatment
 - a was less effective than trolamine for the prevention of skin toxicity
 - b was statistically significantly superior to trolamine for the prevention of skin toxicity
 - c was considered more difficult to apply than trolamine
 - d was associated with several allergic reactions
- 5 According to German Commission E, calendula
 - a flower preparations are suitable for promotion of wound healing
 - b taken internally is effective for the promotion of wound healing
 - c taken internally is effective for GI symptoms such as diarrhoea
 - d flower preparations are suitable for use in inflammation of the oral cavity

CPE ARTICLE 20 ANSWERS: fenugreek

JCM 2005;4(5):82 – SEPTEMBER/OCTOBER 2005

1. d 2. a, b 3. a, b, c 4. b, d 5. a, b

Your details - please print clearly or type

SURNAME		
GIVEN NAMES		
ADDRESS	••••••	
•••••	•••••	••••••
CITY/TOWN	STATE	POSTCODE
TELEPHONE		
EMAIL		

PROGRAM EVALUATION -

Your	r comments on o	our CPE articl	le are appr	eciated		
Rele	vance of this topic					
	very good	good	fair	poor		
Accuracy of the article						
	very good	good	fair	poor		
Education value						
	very good	good	fair	poor		
How many minutes did it take you to complete this unit?						
	< 30 min	30-60	61-90	> 90		
Please tick one						
	ACPPM member – Calendula monograph CPE costs no extra to ACPPM members and subscribers					
	I currently subscribe to the CPE self-assessment program (\$150 for 15 tests)					